No.



8500086

to all to whom these eresents shall come:

Howard J. Hunk & Sons, Inc.

Tolhereas, there has been presented to the

#### Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE; IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EX-LUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, MPORTING IT, OR EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT Y THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

CORN

'DJ7'

In Testimony Wathereof, I have hereunto set my hand and caused the seal of the Blant Tariety Protection Office to be affixed at the City of Washington, D.C.

day of February the year of our Lord one thousand nine hundred and eighty-six

#### Application is required in order to determine WAREHOUSE & SEED DIVISION if a plant variety protection cartificate is to be issued (7 U.S.C. 2421). Information is APPLICATION FOR PLANT VARIETY PROTECTION CERTIFICATE held confidential until certificate is iss (Instructions on reverse) (7 U.S.C. 2426). 1. NAME OF APPLICANT(S) 3. VARIETY NAME 2. TEMPORARY DESIGNATION Edward J. Funk & Sons, Inc. DJ7 4. ADDRESS (Street and No. or R.F.D. No., City, State, and Zip Code) FOR OFFICIAL USE ONLY 5. PHONE (include area code) P.O. Box 67 PVPO NUMBER 601 Funk Parkway 8500086 (219) 474-5111 Kentland, IN 47951 6. GENUS AND SPECIES NAME 7. FAMILY NAME (Botanical) DATE 3<u>/27/8</u>5 Zea mays L. Gramineae TIME 2:30 [ XXP.M. A.M. AMOUNT FOR FILING 8. KIND NAME DATE OF DETERMINATION 800 RECEIVED Yellow dent March 1, 1981 AMOUNT FOR CERTIFICATE 10. IF THE APPLICANT NAMED IS NOT A "PERSON," GIVE FORM OF ORGANIZATION (Corporation, partnership, association, etc.) DATE Corporation 11. IF INCORPORATED, GIVE STATE OF INCORPORATION 12. DATE OF INCORPORATION Indiana 9/5/56 13. NAME AND ADDRESS OF APPLICANT REPRESENTATIVE(S), IF ANY, TO SERVE IN THIS APPLICATION AND RECEIVE ALL PAPERS Dr. Sue Sullivan Edward J. Funk & Sons, Inc. P.O. Box 67 PHONE (include area code): (219) 474-5111 Kentland, IN 47951 14. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED Exhibit C. Objective Description of the Variety (Request form Exhibit A. Origin and Breeding History of the Variety (See from Plant Variety Protection Office.) Section 52 of the Plant Variety Protection Act.) b. X Exhibit D, Additional Description of the Variety Exhibit B, Novelty Statement RIS $\varepsilon$ $\mathbf{X}$ EXHIBIT E, STATEMENT OF BASIS OF OWNERSHIP DOES THE APPLICANTIS) SPECIFY THAT SEED OF THIS VARIETY BE SOLD BY VARIETY NAME ONLY AS A CLASS OF CERTIFIED SEED? (See Section 83(a) of the Plant Variety Protection Act.) Yes (If "Yes," answer items 16 and 17 below) IF "YES" TO ITEM 16, WHICH CLASSES OF PRODUCTION 16. DOES THE APPLICANTIS) SPECIFY THAT THIS VARIETY BE LIMITED AS TO NUMBER OF GENERATIONS? BEYOND BREEDER SEED? Certified Registered 18. DID THE APPLICANT(S) FILE FOR PROTECTION OF THE VARIETY IN THE U.S.? Yas (If "Yes," give date) No N 19 HAS THE VARIETY BEEN OFFERED FOR SALE OR MARKETED IN THE U.S. OR OTHER COUNTRIES? Yes (If "Yes," give names of countries and dates! 20. The applicant(s) declare(s) that a viable sample of basic seeds of this variety will be furnished with the application and will be replenished upon request in accordance with such regulations as may be applicable. The undersigned applicant(s) is (are) the owner(s) of this sexually reproduced novel plant variety, and believe(s) that the variety is distinct, uniform, and stable as required in Section 41, and is entitled to protection under the provisions of Section 42 of the Plant Variety Protection Act. Applicant(s) is (are) informed that false representation herein can jeopardize protection and result in penalties. SIGNATURE OF APPLICANT 3/25/85 SIGNATURE OF APPLICANT

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE

FORM WA-470 (3-84) (Formerly LMGS-470 (8-83) which is obsolete)

FORM APPROVED, OMB NO, 0581-0055

#### EXHIBIT A

#### Origin and Breeding of DJ7

DJ7 was derived from the cross B73xBS16 (Iowa early Eto synthetic) and subsequently backcrossed to B73 three times. Three generations of selfing and ear row selection followed until the uniform DJ7 selection was made in 1981. The first initial increase of the line was made in Hawaii during the winter of 1981-82. To date no known variants have occurred and the line is consistently uniform and stable in its phenotype from year to year.

#### EXHIBIT B

#### Novelty Statement

DJ7 resembles B73 in its phenotypic appearance and hybrid performance. DJ7 differs from B73 in the following ways (see Table 1):

- 1. DJ7 sheds pollen two days earlier than B73.
- 2. DJ7 silks 2.4 days earlier than B73.
- 3. DJ7 is 2.5" shorter than B73.
- 4. DJ7's ear height is three inches shorter than B73.

TABLE 1 Days to Flowering, Plant and Ear Height

	+				Flowe	ring*					· -				
Line	#Reps	50% Pollen			50%	Silk		Plant Ht.(") Ear H		Ht.(")					
		1982	1983	1984	x	1982	1983	1984	x	1982	1984	x	1982	1984	x
В73	38	75.6	76.4	78.0	76.7	78.3	78.5	79.7	78.8	76	<b>7</b> 2	74.0	36	32	34
DJ7	41	73.2	74.3	76.4	74.6	75.2	76.0	78.0	76.4	69	74	71.5	30	32	31

<sup>+</sup> Total number of reps (rows) over years.

<sup>\*</sup> Days from planting to 50% pollen shed and 50% silk.

FORM GR-470-28 (2-15-74)

UNITED STATES DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE GRAIN DIVISION
HYATTSVILLE, MARYLAND 20782

EXHIBIT C (Corn)

### OBJECTIVE DESCRIPTION OF VARIETY

CORN (ZEA MATS)	,
NAME OF APPLICANTIS	FOR OFFICIAL USE ONLY
Edward J. Funk & Sons, Inc.  ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code)	PYPO NUMBER 8500086
P.O. Box 67	VARIETY NAME OR TEMPORARY
601 Funk Parkway	DJ7
Kentland, IN 47951	<i>J</i> J /
Place the appropriate number that describes the varietal character of this variety in the Place a zero in first box (e-s. 0 8 9 or 0 9 ) when number is either 99 or less o	boxes below. r 9 or less.
1. TYPE:	
2 1 = SWEET 2 = DENT 3 = FLINT 4 = FLOUR 5 = F	POP 6 = ORNAMENTAL
2. REGION WHERE BEST ADAPTED IN THE U.S.A.:	
2 1 = NORTHWEST 2 = NORTHCENTRAL 3 = NORTHEAST 5 = SOUTHCENTRAL 6 = SOUTHWEST 7 = MOST REGIONS	
	· omments'' (pg. 3) state how its were calculated)
DAVE SHOW EMERGENCE TO 50% OF PLANTS IN SILK	HEAT UNITS
<u></u>	1810
N A DAYS FROM 50% SILK TO OPTIMUM EDIBLE QUALITY	HEAT UNITS
5 5 DAYS FROM 50% SILK TO HARVEST AT 25% KERNEL MOISTURE 2 5	7 0 HEAT UNITS
4. PLANT:	
1 8 2 CM. HEIGHT (To tassel tip)	CM. EAR HEIGHT (To base of top ear
2 0 CM. LENGTH OF TOP EAR INTERNODE	·
·	
Number of Tillers: Number of Ears Per Stall	k:
To the species	2 = SLIGHT TWO-EAR TENDENCY
2 1 = NONE 2 = 1-2 3 = 2-3 4 = >3 2 1 = SINGLE 3 = STRONG TW	O-EAR TENDENCY 4 THREE-EAR TENDENC
Cytopiasm Type:	
1 = NORMAL 2 = "T" 3 = "S" 4 = "C" 5 = OTHE	R (Specify)
5. LEAF (Field Corn Inbred Examples Giren):	
Color:	
3 1 = LIGHT GREEN (HY) 2 = MEDIUM GREEN (WF9) 3 = DARK GR	REEN (B14) 4 = VERY DARK GREEN (K1
Angle from Stalk (Upper half): Sheath Pubscence:	
1 1 = < 30° 2 = 30-60° 3 => 60° 2 1 = LIGH	T (W22) 2 = MED(UM (WF9)
Marginal Waves: Longitudinal Creases:	
2 1 = NONE (HY) 2 = FEW (WF9) 3 = MANY (OH7L) 2 1 = ABSE 3 = MANY	NT (OH51) 2 = FEW (OH56A) Y (PA11)
Width: Langth:	
1 0 CM WIDEST BOINT OF EAR NODE LEAF 0 7 5 CM.	EAR NODE LEAF
10 CM WIDEST POINT OF EAR NODE LEAF 0 7 5 CM.	
1 3 NUMBER OF LEAVES PER MATURE PLANT	

6.	TASSEL:		8500086
	0 6 NUMBER OF LATERAL BRANCHES		
	Branch Angle from Cantral Spike:	enduncle Length:	
	1 = < 30° 2 = 30-40° 3 = > 45°	CM. FROM TOP LEAF TO	BASAL BRANCHES
	Pollen Shed:		
	1 = LIGHT (WF9) 2 = MEDIUM	3 = HEAVY(KY21)	
	6 Anther Color: 1 = YELLOW 2 = PINK 6 Glume Color: 6 = OTHER (Specify) Anther - Glume - Pollen Restoration for Cytoplasms (o = Not Tested, 1 = Partial, 2 = Good	- purplish green	5 = GREEN
-	0 "т" 0 "s" 0 "c" отн	HER (Specify Cytoplasm and degrees of restoral	tion)
7	7. EAR (Husked Ear Data Except When Stated Otherwise):		
	1 5 CM LENGTH 5 5 MM, MID-POINT DIAMETER	1 2 7 GM, WEIGHT	
	Xernel Rows:  2 1 = INDISTINCT 2 = DISTINCT	1 8 NUMBER	
	1 = STRAIGHT 2 = SLIGHTLY CURVED	3 = SPIRAL	p = 11
	Silk Color (Exposed at Silking Stage):		
	1 = GREEN 2 = PINK 3 = SALMON	4 = RED	·
	Husk Color:		•
	2 FRESH 1 = LIGHT GREEN	2 = DARK GREEN 3 = PINK	
	6 DRY ) 4= RED 5= PUI	RPLE 6 = SUFF	
	Husk Extention: (Harvest Stage)	Husk Leaf:	
٠.	1 = SHORT (Ears Exposed) 2 = MEDIUM (Barely Covering Ear) 3 = LONG (8-10CM Beyond Ear Tip) 4 = VERY LONG (> 10 CM)	2 3 = LONG ( > 15 CM)	MEDIUM (8-15 CM)
	Shank:	Position at Dry Husk Stage:	
	CM LONG NO. OF INTERNODES	1 = UPRIGHT 2 = HORI	ZONTAL 3 = PENDENT
	Taper:	Drying Time (Unhusked Ear):	•
	1 SLIGHT 2 AVERAGE 3 EXTREME	2 1 = SLOW 2 = AVEF	RAGE 3 = FAST
8.	KERNEL (Dried);		
	Size (From Ear Mid-Point):	· · · · · · · · · · · · · · · · · · ·	
	1 3 MM LONG 0 9 MM, WIDE 0	6 MM. THICK	
	Shape Grade (% Rounds)		
	1 = < 20 2 = 20 - 40 3 = 40 - 60	4 = 60 -80 5 = > 80	

÷

1	Pericarp Color:	Color: 1 = COLORLESS 2 = RED-WHITE 5 = BROWN 6 = LIGHT RED 8 = VARIEGATED (Describe)			4 = 8 RRY RED	HONZE			
		8 = VARIEGATE	ED (Describe)	<u> </u>					
1	Aleurone Color:	1 = HOMQZYGO	us	2 = SEG	REGATING (Describ	e)			
1	1 = WHITE	2 = 21NK	1AT = E	ч	4 = BROWN	i	5 = BRC	NZE 6 = RE	
<u> </u>	7 = PURPLE	8 = PALE PU	RPLE	9 = VAF	RIEGATED (Describe	a)			
3	Endosperm Color:	1 = WHITE	2 = PALE Y	ELLOW	3 = YELLOW	. 4.≖PIN	K-ORANGE	5 = WHITE CAP.	
Endosper	m Type:					4			
L		2 = E>	KTRA SWEET (	(sh2)	3 = NORMAL S	TARCH	4 = HIGH AN	YLOSE STARCH	
3		CH 6 = HI			7 = HIGH LYS	NE	8 = OTHER (	(Specify)	
2 5	GM, WEIGHT /10	0 SEEDS (Unsized	Sample)				<u> </u>		
9. COB:									
2 7	MM. DIAMETER	AT MID-POINT			•				
Strength	:			С	olor:				
2	1 = WEAK	2 = STRONG			1 = WHITE 5 = VARIEGA		3 = RED 4		
10. DISEAS	E RESISTANCE (O	= Not Tested, 1 = 5	Susceptible, 2 =	Resistant	):				
0	STALK ROT (Die	olodia)	0 STA	LK ROT (	Fusarium)		STALK RO	OT (Gibberells)	
0				THERNI	EAF BLIGHT	Ī	SMUT		
H					221 0210111				
	SOUTHERN RUS	T .	0 con	IN SMUT		<u> </u>	BACTERIA	KC WIL:	
0	BACTERIAL LEA	AF BLIGHT	0 MAI	ZE DWAR	F MOSAIC	(	) STUNT		
0	OTHER (Specify)					_			
11. INSECT	RESISTANCT (O =	Not Tested, 1 = Su	uscaptible, 2 = f	Resistant):					
0	CORNBORER	0	EARWORM		0	\$APBEET <b>L</b> E	ſ	0 APHID	
0	ROOTWORM (No	orthern)	ROOTWORM	(Western)			_		
0	ROOTWORM (So	uthern) 0	OTHER (Speci	ify)		<u> </u>			
12 VARIET	TIES MOST CLOSE	LY RESEMBLING	THAT SUBMIT	TED FOR	THE CHARACTER	S GIVEN:			
CHARA			VARIETY		CHARAC			VARIETY	
Maturity			B73		Kernel Ty	pe	I	373	
Plant Ty			В73.		Quality (8	dible)	_		
Ear Typ	8		B73	•	Usage			373	
REFER	U.S. Department Corn: Culture, Pr Emerson, R.A., G The Mutants of M Stringfield, G.H.! Butler, D.A. 195	.W. Beadle, and A.C aize, 1968. Crop S Maize Inbred Lines	1970 Avi Publ C. Fraser. A Sun Science Society of Ohio. Ohio	nmary of l of Americ A.E.S. But	npany, Westport, Co Linkaga Studies in Ma La. Madison, Wiscons I. 831. 1959. Inbred Lines — PhD.	sin.	.£.S., Mem. 180.		

#### Additional Information

#### A. Yield

DJ7 is improved over B73 as a seed parent due to higher yield. Table 2 shows DJ7 to be higher yielding and dryer at harvest than B73 in replicated trials. This is verified by company production yield data (Table 3) showing DJ7 to be 10 to 23 bushels/acre higher yielding than B73. The difference is even more pronounced in dry stress years. It is our opinion this is due to the earlier flowering of DJ7 and a tendency for DJ7 silk to remain viable longer under dry conditions.

TABLE 2
Harvest Yield and Moisture

	В73			DJ7			
	1981 <sup>+</sup>	1982*	×	1981+	1982*	x	
Bushels/Acre	64.7	66.0	65.4	75.3	85.1	80.2	
% Moisture	38.8	24.6	31.7	34.8	22.0	28.4	

<sup>+ 2</sup> reps, RCB

<sup>\* 4</sup> reps, RCB

## Additional Information (continued)

#### A. Yield (continued)

TABLE 3

Yield Summary

DJ7 versus B73 - All Pollinators\*

<u>Year</u>	Inbred	Shelled Bu.	Female Acres	Shelled Bu./ Female Acre	DJ7 Advantage Bu./Acre
1983	в73	9,820	1401.0	7.01	
1983	DĴ7	7,916	262.5	30.16	+ 23.15
1984	B73	64,843	1851.0	35.03	. 10 ==
1984	DJ7	48,301	1055.0	45.78	+ 10.75

While the above data depict DJ7's overall superiority in combination with various pollinators, the following data shows how DJ7 performed significantly better with LH38 in 1983 during severe heat and drought stress.

Pedigree	Female Acres	Bushels/ Female Acre	DJ7 Advantage Bu./Acre
DJ7xLH38	8	16.74	. 10.00
B73×LH38	76	2.86	+ 13.88

The above data is from adjacent plantings in 1983. The ability of DJ7 to silk better than B73 under stress conditions accounts for this very significant difference in yield.

<sup>\*</sup>Pollinators for DJ7 were LH38, LH47, LH50, LH51 and LH24. Pollinators for B73 were LH38, LH51, LH24, C123-11, Pa91, L30 and Va22.

## Additional Information (continued)

#### B. Germination Data

#### DJ7xLH38 versus B73xLH38

Based on germination data from the 1984 crop, DJ7 exhibited superior cold germination scores especially in medium round and small round grade sizes. B73 historically shows germination problems in these specific grades.

## \*Cold Germination Data DJ7xLH38 versus B73xLH38

	LR 26/21	LF 26/21		MF 21/17			Plateless
DJ7xLH38	85.20	93.60	91.90	94.50	95.00	95.00	88.00
B73xLH38	84.00	96.00	86.00	95.50	87.00	94.50	86.00

DJ7 average cold germination, 91.4 B73 average cold germination, 89.1

DJ7 shows an overall advantage of 2.3 cold germination points and a 5.9 point advantage in the medium round size plus a 7.0 point advantage in the small round grade.

\*Cold germination is conducted in actual soil medium, 10 days at  $50^{\circ}F$  and 4 days at  $77^{\circ}F$ .

### Additional Information (continued)

#### B. Germination Data (continued)

Warm germination data also indicates an advantage with DJ7, notably in the medium and small round grades.

# \*\* Warm Germination Data DJ7xLH38 versus B73xLH38

	LR 26/21		MR 21/17	MF 21/17	SR 17/15	SF 17/15	Plateless
DJ7xLH38	97.13	97.35	97.81	98.50	96.26	98.00	97.50
B73xLH38	96.75	98.50	95.50	99.25	95.25	97.75	96.50

DJ7 average warm germination, 97.49 B73 average warm germination, 97.03

An overall warm germination advantage of .5 point is realized for DJ7 in all grades. The advantage noted in the medium round size is 2.31 for DJ7 and there is a 1.0 point advantage for DJ7 in the small round grade.

<sup>\*\*</sup> Warm germination is a wet towel test; 7 days at 77°F.

#### EXHIBIT E

Corn Application #8500086; "DJ7"

The corn inbred line, DJ7, is solely owned by Edward J. Funk and Sons, Inc., and all rights to said line are held by the company.